



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Sex and gender differences in COVID-19: an Italian local register-based study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051506
Article Type:	Original research
Date Submitted by the Author:	22-Mar-2021
Complete List of Authors:	Fortunato, Francesca; University of Foggia, Department of Medical and Surgical Sciences; Policlinico Riuniti University Hospital of Foggia, Department of Hygiene Martinelli, Domenico; University of Foggia, Department of Medical and Surgical Sciences; Department of Hygiene Foggia, Policlinico Riuniti University Hospital of Lo Caputo, Sergio; Policlinico Riuniti University Hospital of Foggia, Department of Clinical and Experimental Medicine Santantonio, Teresa; Policlinico Riuniti University Hospital of Foggia, Department of Clinical and Experimental Medicine Dattoli, Vitangelo; Policlinico Riuniti University Hospital of Foggia Lopalco, Pier Luigi; University of Pisa, Department of Translational Research and New Technologies in Medicine and Surgery Prato, Rosa; University of Foggia, Department of Medical and Surgical Sciences
Keywords:	Public health < INFECTIOUS DISEASES, COVID-19, Infection control < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Sex and gender differences in COVID-19: an Italian local register-based study

Francesca Fortunato¹, Domenico Martinelli¹, Sergio Lo Caputo², Teresa Santantonio², Vitangelo Dattoli³, Pier Luigi Lopalco⁴, and Rosa Prato¹.*

¹ Department of Medical and Surgical Sciences, University of Foggia; Department of Hygiene, Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; francesca.fortunato@unifg.it, domenico.martinelli@unifg.it

² Department of Clinical and Experimental Medicine, University of Foggia; Infectious Diseases Unit, Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; sergio.locaputo@unifg.it, teresa.santantonio@unifg.it

³ Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; vitangelodattoli@gmail.com

⁴ Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy; p.lopalco@regione.puglia.it

*Correspondence: rosa.prato@unifg.it; Ospedale "Colonnello D'Avanzo", Viale degli Aviatori, 2 - 71122 Foggia, Italy; Tel.: +39-0881-733091 (R.P.)

Word count: 2778

Keywords: COVID-19; hospitalisation; virus clearance; death; sex differences; gender differences; Italy

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives

The present study aimed to explore differences in COVID-19 outcomes between male and female cases in the Apulian District of Foggia, Italy.

Design and setting

We performed a retrospective epidemiological study among COVID-19 confirmed cases occurred in the Apulian District of Foggia from February 29th to June 30th, 2020. The surveillance data from a regional registry (GIAVA-COVID©) were used.

Main outcomes

The main outcome measures were the proportion of hospitalisations, virus clearance and the case fatality rate.

Results

A total of 1,175 cases (50.7% female; median age: 55 years) were identified. The proportion of hospitalisation with COVID-19 diagnosis was 45.4% in men vs. 37.9% in women ($p<0.01$) while the average length of stay in hospitals was 31.3 ± 14.6 days in women vs. 26.8 ± 14.4 days in men ($p<0.01$). The proportion of cases who achieved virus clearance was higher in women (84.2%; days to clearance: 28.0 ± 12.1) than in men (79.3%; days to clearance: 29.4 ± 12.9 ; $p<0.05$). Men were associated with a significantly higher risk of dying from COVID-19 than women (case fatality rate [CFR] 16.1% vs. 10.4%; $p<0.01$). The mean time, from diagnosis to death was 14.5 ± 14.4 days in women compared with 10.6 ± 10.7 days in men ($p<0.01$). The male sex, age ≥ 55 years and underlying comorbidities significantly raised the risk of hospitalisation, persistent infection and death ($p<0.05$).

Conclusions

This study suggests that more attention should be paid to sex as a variable for the interpretation of COVID-19 data. Sex-disaggregated data will help the clinicians to make appropriate patient-tailored medical decisions.

Strengths and Limitations of this Study

- This study provides sex-disaggregated data of COVID-19 cases at a district level, in Italy. These data can contribute to a better understanding of who is being impacted the most by the pandemic and promote a patient-tailored treatment approach.
- The robust methodology of the present study enabled to accurately correlate the case demographics with COVID-19 clinical response. In this context, data related to the viral clearance, which reflect the diversified course of the disease according to the individual immune response, are confirmatory of sex difference in COVID-19.
- The data collected are highly homogeneous as they are strictly related to the first epidemic wave and provide an accurate picture of the impact of sex and age on COVID-19 outcomes in Italy during the initial phase of the pandemic.
- As the majority of the sex-disaggregated data available in the literature, the data presented in our study are not adjusted for lifestyle, profession, social or behavioural differences.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel coronavirus causing the current pandemic, which has resulted in millions of infections and hundreds of thousands of deaths worldwide. As of March 10th 2021, a total of 3,069,625 cases of confirmed SARS-COV-2 have been reported in Italy with a case fatality rate (CFR) of 3.2%.^[1]

The clinical manifestations of SARS-CoV-2 vary from asymptomatic infection to severe or critical disease.^[2]

Older age and comorbidities such as hypertension, cardiovascular disease, diabetes, chronic respiratory disease are associated with severe disease and death.^[3–5] Sex and gender have been identified as additional risk factors contributing to heterogeneous COVID-19 outcomes.^[2] Indeed, several studies have reported sex bias in COVID-19 case fatalities. It is observed that men have a higher risk of developing a severe form of the disease compared with women, highlighting the importance of sex-disaggregated data of COVID-19 cases.^[6] The initial reports from China followed by data from several European countries have shown similar numbers of confirmed cases between men and women.^[7,8] However, the severity of COVID-19, measured as hospitalisation, admission to intensive care units and fatality rate, is 2-fold higher for men than women.^[8,9] Studies in China, South Korea, United States, United Kingdom and Italy have reported higher case fatality rates and worst disease outcomes in male cases than in female cases.^[7,10–14] In some of these studies, the higher fatality rate in men was observed even after adjusting for confounding factors such as age and comorbidities.^[7,14] Additionally, in Italy, the higher fatality rate in men (age range: 40-80 years) is confirmed when the healthcare worker population is selectively studied.^[1]

The reasons for the differences in COVID-19 outcome and progression between men and women remain unclear. On one hand, biological factors, such as chromosomal and hormonal differences between men and women, may influence their susceptibility to infections, immunologic responses and progression of the disease.^[9,15–17] On the other hand, gender-related factors including psychological, social and behavioural differences between men and women may affect SARS-CoV-2 exposure, presence of comorbidities, treatment initiation and compliance, and COVID-19 mortality.^[18,19]

In this study, we used the surveillance data from a regional registry containing all confirmed cases of COVID-19 occurred in the Foggia District (Apulia region, Italy), as of late June 2020, after the end

of the first epidemic wave. We aimed to explore the sex differences in hospitalisation, virus clearance, and deaths.

METHODS

Study population and design

We conducted a retrospective epidemiological study among COVID-19 cases occurred in the Foggia District, Apulia region, Italy, from February 29th to June 30th, 2020. Foggia District is the third-largest Apulian District, with an estimated 616,310 residents (51% women) as of January 1st 2020.[20]

We used the surveillance data from a regional registry (GIAVA – COVID©) developed based on the Go.Data outbreak investigation tool (WHO) to manage the emergency.[21] GIAVA – COVID© includes functionalities for investigation and follow-up of cases and contacts, contact tracing, laboratory and clinical data collection. Information collected include age, sex, residence location, date of illness onset, date of diagnosis, date of hospital admission, date of COVID-19 positive and negative tests, date of death, presence of underlying diseases, case outcomes (hospitalisation, virus clearance and death), and disease severity (mild, moderate, severe, or critical).[22] The disease classification is duly updated according to the change in clinical manifestations of each case.

This study included laboratory-confirmed cases defined as any person meeting the laboratory criterion (detection of SARS-CoV-2 nucleic acid or antigen in a clinical specimen).[23]

The proportion of hospitalisation was defined as the proportion of all infected individuals who were hospitalised among the total number of infected individuals. The proportion of individuals who achieved virus clearance was defined as the proportion of those clinically recovered and who had laboratory evidence of viral RNA clearance from the upper respiratory tract among the total number of infected individuals. The case fatality rate was defined as the proportion of deaths from diagnosed cases among the total number of infected individuals.

Statistical Analysis

Categorical variables were summarised as the counts and percentages in each category. Continuous variables were expressed as the medians and interquartile ranges (IQR) and the means (\pm standard deviation [SD]). Differences in continuous variables were tested with Student's t-test for normally distributed ones, or the Mann-Whitney U test when variables showed a non-normal distribution.

1
2
3 121 Significant differences in categorical variables were assessed using the Chi-square test or Fisher’s
4
5 122 exact test when appropriate. Multivariate logistic regression analysis was performed to evaluate
6
7 123 whether demographics (sex: male vs. female; age group: above the median age vs. below the
8
9 124 median age) and clinical characteristics were independently associated with hospitalisation, virus
10
11 125 clearance, and deaths. The analysis was conducted with STATA/SE 15.0.
12
13

14 126 **Patient and public involvement**

15
16 127 Patients and/or the public were not involved either in the study design, conduct, reporting or in the
17
18 128 dissemination plans of this research.
19
20 129

23 130 **RESULTS**

25
26 131 Between February 29th and June 30th, 2020, a total of 1,175 cases (50.7% female; median age: 55
27
28 132 years, IQR: 40-71 years) were diagnosed with COVID-19 in the Foggia District, Apulia region, Italy.
29
30 133 Comparison of demographics and clinical characteristics of men versus women are shown in Table
31
32 134 1. A total of 373 cases (31.7%) had underlying medical conditions, including cardiovascular disease
33
34 135 (63.3%), diabetes (19.6%), chronic pulmonary disease (13.9%), cancer (10.7%), neurological diseases
35
36 136 (9.9%), chronic kidney disease (9.4%), and obesity (with BMI between 30-40 kg/m² or higher) (6.7%).
37
38 137 Nearly 50% of cases were asymptomatic or with mild disease, 14.4% had moderate disease, 20.9%
39
40 138 developed a severe disease and 3.2% progressed to a critical stage. There was no significant
41
42 139 difference in age, underlying comorbidities (with the exception of diabetes), and disease severity
43
44 140 distributions between the male and female groups (Table 1).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Comparison of characteristics between male and female COVID-19 cases in the Foggia District, Apulia region, Italy

Characteristics	Male	Female	Total	OR (95% CI)	χ^2	p value
No. of cases (%)	579 (49.3)	596 (50.7)	1,175			
Median age (IQR), years	56 (40-70)	54.5 (38-74)	55 (40-71)			
Mean age (\pmSD), years	54.3 \pm 21.1	54.5 \pm 22.6	54.4 \pm 21.8			0.4291
Age group, No. (%):						
0-9	14 (2.4)	16 (2.7)	30 (2.6)	Ref.		
10-19	23 (4.0)	22 (3.7)	45 (3.8)	1.19 (0.43-3.34)	0.1	0.7061
20-29	43 (7.4)	53 (8.9)	96 (8.2)	0.93 (0.38-2.30)	0.03	0.8571
30-39	57 (9.8)	64 (10.7)	121 (10.3)	1.02 (0.42-2.47)	0.00	0.9655
40-49	91 (15.7)	95 (15.9)	186 (15.8)	1.09 (0.47-2.57)	0.05	0.8184
50-59	105 (18.1)	108 (18.1)	213 (18.1)	1.11 (0.48-2.59)	0.07	0.7874
60-69	96 (16.6)	64 (10.7)	160 (13.6)	1.71 (0.72-4.07)	1.84	0.1747
70-79	71 (12.3)	69 (11.6)	140 (11.9)	1.17 (0.49-2.81)	0.16	0.6874
80-89	64 (11.1)	72 (12.1)	136 (11.6)	1.01 (0.43-2.44)	0.00	0.9689
≥ 90	15 (2.6)	33 (5.5)	48 (4.1)	0.52 (0.18-1.48)	1.88	0.1705
Comorbidity, No (%)	191 (33.0)	182 (30.5)	373 (31.7)	1.1 (0.86-1.43)	0.75	0.3860
Cardiovascular disease	126 (66.9)	110 (60.4)	236 (63.3)	1.27 (0.81-1.97)	1.23	0.2682
Diabetes	49 (25.7)	24 (13.2)	73 (19.6)	2.27 (1.29-4.01)	9.20	0.0024
Chronic pulmonary disease	30 (15.7)	22 (12.1)	52 (13.9)	1.35 (0.72-2.58)	1.02	0.3132
Cancer	23 (12.0)	17 (9.3)	40 (10.7)	1.32 (0.65-2.75)	0.71	0.3994
Neurological diseases	15 (7.9)	22 (12.1)	37 (9.9)	0.62 (0.29-1.30)	1.87	0.1715
Chronic kidney disease	22 (11.5)	13 (7.1)	35 (9.4)	1.69 (0.78-3.78)	2.10	0.1475
Obesity	13 (6.8)	12 (6.6)	25 (6.7)	1.03 (0.42-2.55)	0.01	0.9345
Other metabolic diseases	5 (2.6)	10 (5.5)	15 (4.0)	0.46 (0.12-1.52)	2.00	0.1575
Liver disease	10 (5.2)	4 (2.2)	14 (3.8)	2.45 (0.69-10.91)	2.38	0.1228
Disease Severity, No. (%)						
Critical	23 (4.0)	15 (2.5)	38 (3.2)	1.88 (0.88-4.11)	3.15	0.0760
Severe	126 (21.8)	120 (20.1)	246 (20.9)	1.28 (0.87-1.90)	1.80	0.1796
Moderate	80 (13.8)	89 (14.9)	169 (14.4)	1.10 (0.72-1.69)	0.23	0.6348
Mild	165 (28.5)	197 (33.1)	362 (30.8)	1.02 (0.72-1.47)	0.03	0.8718
Asymptomatic	92 (15.9)	113 (19.0)	205 (17.4)	Ref.		

CI: confidence interval; IQR: interquartile range; OR: odds ratio; Ref.: reference group; SD: standard deviation

1
2
3 145 The proportion of hospitalisation with COVID-19 diagnosis was estimated to be 41.6%, with a
4
5 146 significant difference observed between men (45.4%) and women (37.9%; $p<0.01$). While the
6
7 147 average length of stay in hospitals was significantly higher in women (31.3 ± 14.6 days) than in men
8
9 148 (26.8 ± 14.4 days; $p<0.01$), there were more women aged ≥ 55 years hospitalised ($p<0.01$). The
10
11 149 proportion of cases who achieved virus clearance was 82%, higher in women (84.2%; days to
12
13 150 clearance: 28.0 ± 12.1) than in men (79.3%; days to clearance: 29.4 ± 12.9 ; $p<0.05$). A total of 155
14
15 151 deaths occurred among all cases for an overall CFR of 13.2%. Men were associated with a
16
17 152 significantly higher risk of dying from COVID-19 than women (16.1% vs. 10.4%; $p<0.01$). The mean
18
19 153 time, from diagnosis to death was higher in women (14.5 ± 14.4 days) compared with men (10.6 ± 10.7
20
21 154 days; $p<0.01$) (Table 2).
22
23 155

24 156 **Table 2. Comparison of outcomes between male and female COVID-19 cases in the Foggia District, Apulia**
25 157 **region, Italy**

Characteristics	Male	Female	OR (95% CI)	χ^2	p value
Hospitalisation, No. (%)	263 (45.4)	226 (37.9)	1.36 (1.07-1.73)	6.81	0.0091
Mean age (\pm SD), years	66.2 \pm 16.0	70.2 \pm 18.8			0.0053
Mean length-of-stay in hospital (\pm SD), days	26.8 \pm 14.4	31.3 \pm 14.6			0.0032
Mean length-of-stay in hospital (IQR), days	24 (17-35)	29 (19-41)			
Virus clearance (yes), No. (%)	459 (79.3)	502 (84.2)	0.72 (0.53-0.97)	4.84	0.0278
Mean time-to-virus-clearance (\pm SD), days	29.4 \pm 12.9	28.0 \pm 12.1			0.0432
Median time-to-virus-clearance (IQR), days	25 (18-35)	27 (19-37)			
Deaths, No. (%)	93 (16.1)	62 (10.4)	1.65 (1.15-2.36)	8.21	0.0042
Mean time-to-death (\pm SD), days	10.6 \pm 10.7	14.5 \pm 14.4			0.0282
Median time-to-death (IQR), days	8 (3-16)	10 (4-23)			

48 158 CI: confidence interval; IQR: interquartile range; OR: odds ratio; SD: standard deviation
49

50 159
51
52 160 The male sex, age ≥ 55 years and underlying comorbidities significantly raised the risk of
53
54 161 hospitalisation, persistent infection and death ($p<0.05$; Table 3).
55
56
57
58
59
60

Table 3. Multivariate logistic regression analysis of demographics and clinical characteristics of COVID-19 cases in the Foggia District, Apulia region, Italy

Characteristics	Hospitalisation		Virus clearance (no)		Deaths	
	OR (95% CI)	p value	OR (95% CI)	p value	OR (95% CI)	p value
Sex (male vs. female)	1.52 (1.15-2.20)	0.003	1.51 (1.08-2.09)	0.014	2.33 (1.52-3.58)	0.000
Age group (≥55 years vs. <55 years)	1.83 (1.68-1.99)	0.000	1.62 (1.47-1.78)	0.000	2.62 (2.22-3.07)	0.000
Comorbidity (yes vs. no)	1.99 (1.47-2.69)	0.000	1.63 (1.16-2.29)	0.004	1.94 (1.28-2.93)	0.002

CI: confidence interval; OR: odds ratio

DISCUSSION

Our registry-based surveillance study of 1,175 COVID-19 cases, well characterised from both demographic and clinical points of view, highlighted a male bias in COVID-19 outcomes. Based on the herein presented data, men are more likely to be hospitalised compared with women and the proportion of male cases achieving virus clearance is lower compared with female cases. Furthermore, men require longer periods to achieve virus clearance, have a higher fatality rate and faster progression to death.

A male bias (male-to-female ratio >1.1) in COVID-19 mortality is currently reported in 75 of the 94 countries that have provided sex-disaggregated data (as of March 10th 2021). At the global level, a higher number of men are hospitalised or admitted to the intensive care unit (ICU) compared with women.[24] Additionally, several studies have demonstrated that men with COVID-19 are at higher risk of death and severe form of infection than women.[25,26] A recent meta-analysis of 3,111,714 reported global cases demonstrated that, whilst there is no difference in the proportion of male and female COVID-19 cases, men have higher odds of death (odds ratio [OR]=1.39; 95% confidence interval [CI]=1.31,1.47) compared with women.[27]. Similarly, our study presents a comparable proportion of women and men with confirmed COVID-19 (50.7% vs. 49.3%). Therefore, the observed differences cannot be attributed to a prevalence of COVID-19 in male sex.

The sex distribution of confirmed cases observed in the Foggia District, Apulia region is in line with the overall sex distribution of cases observed in Italy and other European countries.[1,8,28] Although in the early phase of the pandemic in Italy a higher prevalence of COVID-19 was observed in men compared with women, this disproportion became less evident with the progression of the

1
2
3 187 pandemic. This variability may be explained by the different surveillance approaches adopted during
4
5 188 the pandemic since during the first epidemic wave a symptom-based screening led to an
6
7 189 underestimation of asymptomatic to mild cases. In Italy, after the end of the first epidemic wave
8
9 190 (June 30th 2020), a higher number of male cases was observed in the 0-9, 10-19, 60-69 and 70-79
10
11 191 years age groups (52.7%, 50.1%, 59.5%, 57.1%, respectively) compared with female cases, whereas
12
13 192 a nearly 4-times higher number of female cases was observed in the >90 years age group.[29] On
14
15 193 the contrary, as of March 10th 2021, the number of confirmed COVID-19 cases is slightly higher in
16
17 194 women both in the overall Italian territory (51.4% in women vs. 48.6% in men) and in Apulia (51.%
18
19 195 in women vs. 48.4% in men).[1,30]
20
21 196 Differences in disease incidence, morbidity and mortality between sexes have also been observed
22
23 197 in other infectious diseases such as the severe acute respiratory syndrome coronavirus (SARS-COV-
24
25 198 1) and the Middle East respiratory syndrome coronavirus (MERS-COV) with men being more
26
27 199 susceptible than women to the infection and having worse outcome.[31,32] The difference in
28
29 200 mortality between men and women suggests that women are either less prone to develop severe
30
31 201 complications or that they are less likely to die because of severe complications.[33]
32
33 202 The reasons behind these sex-related differences are probably pathogen-specific and of
34
35 203 multifactorial origin.[26] The three main determinants so far proposed to explain male-female
36
37 204 disparities in SARS-CoV-2 infection are differences in immune function associated with the X
38
39 205 chromosome, the effects of sex hormones, gender-related behavioural and socio-cultural
40
41 206 differences.[2,6,16,17] For example, the localisation of angiotensin-converting enzyme-2 (ACE2)
42
43 207 and Toll-like receptor 7 (TLR7) genes to the X chromosome and the mono-allelic versus the bi-allelic
44
45 208 presence may help explain the increased risk of COVID-19 for males compared to females.[34]
46
47 209 From a biological point of view, women seem to have a stronger immune system, weaker cytokine-
48
49 210 based pro-inflammatory response and lower levels of angiotensin-converting enzyme-2 (ACE2), an
50
51 211 essential component for the entrance of COVID-19 into the cells.[2,16,35–37] In this context,
52
53 212 oestrogens seem to play a key protective role. Oestrogen levels vary with age, rising in prepubertal
54
55 213 individuals and decreasing with age. Thus, the age-associated decline in oestradiol levels might be
56
57 214 an explanation for the higher susceptibility and severe progression of COVID-19 in older
58
59 215 subjects.[38]
60
61 216 Our study highlights that, alongside sex, age and comorbidity are risk factors increasing
62
63 217 hospitalisation and death, and decreasing virus clearance. That COVID-19 severity increases with

age became evident since the beginning of the pandemic. Early studies from China and Italy showed that older age was associated with a greater risk of developing acute respiratory distress syndrome (ARDS), severe lung disease and death.[5,10] A recent meta-analysis of 55 studies and 10,014 COVID-19 cases confirmed that older age (≥ 50 years), together with comorbidities, significantly affects the prognosis and severity of COVID-19.[3] A further study investigated whether male bias in COVID-19 mortality was maintained at every age. It analysed data collated by the National Institute for Demographic Studies from national statistical agencies across England and Wales, France, Germany, Italy, Netherlands, Portugal, Korea, and Spain, including a population of 194,349,591 men and 201,715,364 women from the beginning of the pandemic until June 21st 2020. The overall male-to-female mortality sex ratio per 100,000 population was 1.4 (crude ratio 1.3). This ratio varied with age: 0.81 for subjects aged 0-9 years; 1.9 in the 40-49 year age group, 2.3 in the 50-59 year age group; 2.6 in the 60-69 year age group and 1.65 in subjects older than 80 years.[39] How the male vs. female difference in mortality, hospitalisation and virus clearance progresses with age is an aspect that warrants further investigation. In this context, stratification of the sex-disaggregated data provided in our study by age group could be relevant to better understand to what extent women are genetically protected from COVID-19. Interestingly, in our study, the stratification of the population by a cut-off age of 55 years highlighted a higher hospitalisation rate in the subgroup of women aged ≥ 55 years, suggesting the role of the reduction of hormonal protection with age.

One of the main hypotheses that have been postulated to justify the observed sex heterogeneity in the immune response to SARS-CoV-2 infection is the different genetic profile. Increasing evidence from patient populations highlights a substantial contribution of human genetic factors to the diversified susceptibility to SARS-CoV-2 infection and/or COVID-19 severity. In this context, a differential response to COVID-19 has also been observed among individuals with ethnicity-based differences in their genetic profile.[34] For instance, the distribution of the gene cluster on chromosome 3, that has been recently identified as the major genetic risk factor for severe COVID-19, differs among populations of different ethnic background (i.e. Asian, European and African populations).[40]

Lastly, gender-related differences in lifestyles and social roles require careful considerations as they are believed to greatly influence the onset, course and outcome of COVID-19. It has been proposed that smoking and alcohol consumption, alongside poor eating habits, more frequently found in men than women, may lead to a higher incidence of comorbidities in men compared with women

1
2
3 250 explaining the higher male mortality observed on a global level.[18,41] However, it must be noted
4
5 251 that no significant difference in underlying comorbidities (with the exception of diabetes) between
6
7 252 men and women was found in our study. There may be other behavioural and social differences
8
9 253 favouring women as men are more reluctant to follow hand hygiene and seek preventive care.[42]
10
11 254 On the other hand, women might be more easily exposed to SARS-CoV-2 infection in both
12
13 255 professional and household settings. Indeed, women represent 70% of the health and social care
14
15 256 workforce and more often care for household members with COVID-19.[18,41]
16
17 257 The present study aimed to explore the differences in hospitalisation and death between men and
18
19 258 women at the local level taking into consideration COVID-19 confirmed cases in the Apulian District
20
21 259 of Foggia. The results are in line with what observed on a national and global level. Hospitalisation
22
23 260 and death are hard outcomes for monitoring the course and severity of the disease. Furthermore,
24
25 261 sex difference in virus clearance represents an added-value outcome of our study as it expresses
26
27 262 the immune response of the host.
28
29 263 However, it should not be neglected that one of the main limitations of our study is that the
30
31 264 presented data are not adjusted for lifestyle, profession, social or behavioural differences, all
32
33 265 relevant factors that could change the interpretation of the data and could further emphasise the
34
35 266 male-bias in COVID-19 severity and fatality. This limitation is a common feature of the majority of
36
37 267 sex-disaggregated data currently available. Indeed, due to practicability and ethical reasons, no
38
39 268 prospective study comparing an equal number of men and women under equal conditions of viral
40
41 269 exposure has been conducted to date. Therefore, we highlight the need of taking into account the
42
43 270 social, familiar and professional roles, alongside biological variables, in order to fully understand the
44
45 271 differences in COVID-19 outcome between men and women.
46
47 272 The main strength of our study consists in its robust methodology, which enabled an accurate
48
49 273 evaluation of the correlation between the case demographics (especially gender) and COVID-19
50
51 274 clinical response. Specifically, the collection of viral clearance data highlights a statistically
52
53 275 significant male-to-female difference and provides a plausible explanation for the observed
54
55 276 diversified course of the disease. Furthermore, the data collected in our study are highly
56
57 277 homogeneous as they are strictly related to the first epidemic wave, and provide an accurate picture
58
59 278 of the impact of sex and age on SARS-CoV-2 infection response in Italy during the initial phase of the
60
280 pandemic. The ethnical composition of the population included in our study is also highly
281 homogeneous and likely well representative of the Italian population or other Mediterranean
European populations.

CONCLUSIONS

Despite a comparable incidence of COVID-19 among the two sexes, a male bias in COVID-19 mortality is observed in the majority of the countries with available sex-disaggregated data. Our study provides sex-disaggregated data for the COVID-19 cases of the Apulian district of Foggia, Italy. It demonstrates that male sex, alongside older age (age ≥ 55 years) and comorbidity, is associated with a greater risk of hospitalisation and death, and lower virus clearance. Therefore, more attention should be paid to sex as a variable for the interpretation of COVID-19 data. This study will help the clinicians to make appropriate patient-tailored medical decisions based on patient sex, age and comorbidities. Future investigations providing data adjusted for gender-related factors (social, familiar and professional roles) are warranted.

Acknowledgements The Authors would like to thank all the frontline health workers at Hospitals and Local Health Unit for their dedication and valuable work into pandemic control. A special thanks to Lucia Massi, Maria Rosa Valetto and Pietro Dri (Zadig, Scientific Publisher, Milan, Italy) for editorial assistance, manuscript development and writing support.

Contributors FF and RP conceptualised and designed the work, analysed and interpreted data, and wrote the manuscript. DM and PLL supervised the study, coordinated regional data collection and provided statistical support. SLC, TS and VD interpreted the results and critically reviewed the advanced version of the manuscript. All authors approved the final draft of the manuscript. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for profit sectors.

Competing interests The authors declare no competing interests.

Patient and public involvement Patients and/or the public were not involved either in the study design, conduct, reporting or in the dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval

The study was conducted in accordance with the principles expressed in the Declaration of Helsinki 1975, as revised in 2008. As this study constituted public health surveillance, ethical approval from institutional review board was not required. All data were provided and analysed anonymously.

Data availability statement No additional data available.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

For peer review only

REFERENCES

- 1 COVID-19 integrated surveillance data in Italy.
<https://www.epicentro.iss.it/en/coronavirus/sars-cov-2-dashboard> (accessed 10 March 2021).
- 2 Falahi S, Kenarkoohi A. Sex and gender differences in the outcome of patients with COVID-19. *J Med Virol* 2021;**93**:151–2.
- 3 Barek MA, Aziz MA, Islam MS. Impact of age, sex, comorbidities and clinical symptoms on the severity of COVID-19 cases: A meta-analysis with 55 studies and 10014 cases. *Heliyon* 2020;**6**:e05684.
- 4 Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA* 2020;**18**:1775–6.
- 5 Wu C, Chen X, Cai Y, *et al.* Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med* 2020;**180**:934.
- 6 Dana PM, Sadoughi F, Hallajzadeh J, *et al.* An Insight into the Sex Differences in COVID-19 Patients: What are the Possible Causes? *Prehosp Disaster Med* 2020;**4**:438–41.
- 7 Chen J, Bai H, Liu J, *et al.* Distinct Clinical Characteristics and Risk Factors for Mortality in Female Inpatients With Coronavirus Disease 2019 (COVID-19): A Sex-stratified, Large-scale Cohort Study in Wuhan, China. *Clin Infect Dis* 2020;**71**:3188–95.
- 8 Gebhard C, Regitz-Zagrosek V, Neuhauser HK, *et al.* Impact of sex and gender on COVID-19 outcomes in Europe. *Biol Sex Differ* 2020;**11**:29.
- 9 Klein SL, Dhakal S, Ursin RL, *et al.* Biological sex impacts COVID-19 outcomes. *PLOS Pathog* 2020;**16**:e1008570.
- 10 Borghesi A, Zigliani A, Masciullo R, *et al.* Radiographic severity index in COVID-19 pneumonia: relationship to age and sex in 783 Italian patients. *Radiol Med* 2020;**125**:461–4.
- 11 Chen T, Wu D, Chen H, *et al.* Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;**368**:m1091.
- 12 Dudley JP, Lee NT. Disparities in Age-Specific Morbidity and Mortality from SARS-CoV-2 in China and the Republic of Korea. *Clin Infect Dis* 2020;**71**:863–5.
- 13 Gavin W, Campbell E, Zaidi S-A, *et al.* Clinical characteristics, outcomes and prognosticators in adult patients hospitalized with COVID-19. *Am J Infect Control* 2021;**49**:158–65.
- 14 Williamson EJ, Walker AJ, Bhaskaran K, *et al.* Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;**584**:430–6.
- 15 Dana PM, Sadoughi F, Hallajzadeh J, *et al.* An Insight into the Sex Differences in COVID-19 Patients: What are the Possible Causes? *Prehospital and Disaster Medicine*. 2020;**35**:438–41.

1
2
3 352 16 Bienvenu LA, Noonan J, Wang X, *et al.* Higher mortality of COVID-19 in males: sex differences
4 353 in immune response and cardiovascular comorbidities. *Cardiovasc Res* 2020;**116**:2197–206.
5
6
7 354 17 Scully EP, Haverfield J, Ursin RL, *et al.* Considering how biological sex impacts immune
8 355 responses and COVID-19 outcomes. *Nat Rev Immunol* 2020;**20**:442–7.
9
10 356 18 Cataldo Claudia, Masella Roberta. Gender-related sociocultural differences and COVID-19:
11 357 what influence on the effects of the pandemic? *Epidemiol Prev* 2020;**44**:Suppl 2:398-399.
12
13 358 19 Griffith DM, Sharma G, Holliday CS, *et al.* Peer Reviewed: Men and COVID-19: A
14 359 Biopsychosocial Approach to Understanding Sex Differences in Mortality and
15 360 Recommendations for Practice and Policy Interventions. *Prev Chronic Dis* 2020;**17**:E63.
16 360
17
18 361 20 Statistiche demografiche ISTAT. <http://demo.istat.it/pop2020/index.html> (accessed 1 Feb
19 362 2021).
20
21 363 21 Go.Data: Managing complex data in outbreaks. <https://www.who.int/godata> (accessed 1 Feb
22 364 2021).
23 364
24
25 365 22 Clinical management of COVID-19. [https://www.who.int/publications/i/item/clinical-](https://www.who.int/publications/i/item/clinical-management-of-covid-19)
26 366 [management-of-covid-19](https://www.who.int/publications/i/item/clinical-management-of-covid-19) (accessed 10 March 2021).
27 366
28 367 23 Case definition for coronavirus disease 2019 (COVID-19), as of 3 December 2020.
29 368 <https://www.ecdc.europa.eu/en/covid-19/surveillance/case-definition> (accessed 1 Feb 2021).
30 368
31
32 369 24 Global Health 50/50. [https://globalhealth5050.org/the-sex-gender-and-covid-19-](https://globalhealth5050.org/the-sex-gender-and-covid-19-project/dataset/)
33 370 [project/dataset/](https://globalhealth5050.org/the-sex-gender-and-covid-19-project/dataset/) (accessed 10 March 2021).
34 370
35 371 25 Kragholm K, Andersen MP, Gerds TA, *et al.* Association between male sex and outcomes of
36 372 Coronavirus Disease 2019 (Covid-19) – a Danish nationwide, register-based study. *Clin Infect*
37 373 *Dis*;ciaa**924**.
38 373
39
40 374 26 Lakbar I, Luque-Paz D, Mege J-L, *et al.* COVID-19 gender susceptibility and outcomes: A
41 375 systematic review. *PLOS ONE* 2020;**15**:e0241827.
42 375
43 376 27 Peckham H, Gruijter NM de, Raine C, *et al.* Male sex identified by global COVID-19 meta-
44 377 analysis as a risk factor for death and ITU admission. *Nat Commun* 2020;**11**:1–10.
45 377
46
47 378 28 European Centre for Disease Prevention and Control (ECDC). COVID-19 situation board.
48 379 [https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-](https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-transmission-tab)
49 380 [transmission-tab](https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-transmission-tab) (accessed 10 March 2021).
50 380
51
52 381 29 Bollettino-sorveglianza-integrata-COVID-19_30 giugno 2020.
53 382 [https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-](https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_30-giugno-2021.pdf)
54 383 [19_30-giugno-2021.pdf](https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_30-giugno-2021.pdf) (accessed 10 March 2021).
55 383
56 384 30 Bollettino Epidemiologico Regione Puglia. Epidemia COVID-19, 10 marzo 2021.
57 385 https://www.regione.puglia.it/documents/65725/216593/Bollettino+Covid_10032021.pdf/6cf
58 386 [c1d87-a820-25c9-9832-2585c0f50099?t=1615382755715](https://www.regione.puglia.it/documents/65725/216593/Bollettino+Covid_10032021.pdf/6cf) (accessed 10 March 2021).
59 386
60

- 1
2
3 387 31 Karlberg J, Chong DSY, Lai WYY. Do men have a higher case fatality rate of severe acute
4 388 respiratory syndrome than women do? *Am J Epidemiol* 2004;**159**:229–31.
- 6
7 389 32 Alghamdi IG, Hussain II, Almalki SS, *et al.* The pattern of Middle East respiratory syndrome
8 390 coronavirus in Saudi Arabia: a descriptive epidemiological analysis of data from the Saudi
9 391 Ministry of Health. *Int J Gen Med* 2014;**7**:417–23.
- 11 392 33 Penna C, Mercurio V, Tocchetti CG, *et al.* Sex-related differences in COVID-19 lethality. *Br J*
12 393 *Pharmacol* 2020;**177**:4375–85.
- 14
15 394 34 Anastassopoulou C, Gkizarioti Z, Patrinos GP, *et al.* Human genetic factors associated with
16 395 susceptibility to SARS-CoV-2 infection and COVID-19 disease severity. *Human Genomics*
17 396 2020;**14**:40.
- 19 397 35 Haitao T, Vermunt JV, Abeykoon J, *et al.* COVID-19 and Sex Differences: Mechanisms and
20 398 Biomarkers. *Mayo Clin Proc* 2020;**95**:2189–203.
- 22
23 399 36 Takahashi T, Ellingson MK, Wong P, *et al.* Sex differences in immune responses that underlie
24 400 COVID-19 disease outcomes. *Nature* 2020;**588**:315–20.
- 26 401 37 Takahashi T, Iwasaki A. Sex differences in immune responses. *Science* 2021;**371**:347–8.
- 28
29 402 38 Liang X. Is COVID-19 more severe in older men? *Postgrad Med J* 2020;**96**:426.
- 30
31 403 39 Bhopal SS, Bhopal R. Sex differential in COVID-19 mortality varies markedly by age. *The Lancet*
32 404 2020;**396**:532–3.
- 34 405 40 Zeberg H, Pääbo S. The major genetic risk factor for severe COVID-19 is inherited from
35 406 Neanderthals. *Nature* 2020;**587**:610–2.
- 37
38 407 41 Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. *The Lancet*
39 408 2020;**395**:846–8.
- 41 409 42 Sharma G, Volgman AS, Michos ED. Sex Differences in Mortality From COVID-19 Pandemic: Are
42 410 Men Vulnerable and Women Protected? *JACC Case Rep* 2020;**2**:1407–10.
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—checklist of items that should be included in reports of observational studies

Sex and gender differences in COVID-19: an Italian local register-based study
Francesca Fortunato, Domenico Martinelli, Sergio Lo Caputo, Teresa Santantonio, Vitangelo Dattoli, Pier Luigi Lopalco, and Rosa Prato.

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1 (title) 2 (abstract)
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2 (abstract)
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4 (lines 83-89)
Objectives	3	State specific objectives, including any prespecified hypotheses	5 (lines 92,93)
Methods			
Study design	4	Present key elements of study design early in the paper	5 (lines 97-99)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5 (lines 100-107)
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	(a) Cross-sectional 5 (lines 108,09)
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5 (lines 103-107)
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5 (lines 100,101; 110-115)
Bias	9	Describe any efforts to address potential sources of bias	5 (lines 144-147)
Study size	10	Explain how the study size was arrived at	5 (lines 97-99)
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	5 (lines 110-115)

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	5,6 (lines 116-125)
		(b) Describe any methods used to examine subgroups and interactions	6 (lines 122-125)
		(c) Explain how missing data were addressed	Not applicable
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	
		(e) Describe any sensitivity analyses	Not applicable
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6 (lines 132,133) Table 1
		(b) Give reasons for non-participation at each stage	Not applicable
		(c) Consider use of a flow diagram	Not applicable
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6 (lines 132-140) Table 1
		(b) Indicate number of participants with missing data for each variable of interest	Not applicable
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	7-9 (lines 145-151) Tables 2,3
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8,9 (lines 151-161) Tables 2,3
		(b) Report category boundaries when continuous variables were categorized	Tables 2,3
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	8,9 (lines 160-161) Tables 2,3
Discussion			
Key results	18	Summarise key results with reference to study objectives	9 (lines 167-172)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12 (lines 263-271)
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12 (lines 283-281)
Generalisability	21	Discuss the generalisability (external validity) of the study results	12 (lines 272-281)
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	13 (lines 302,302)

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Sex and gender differences in COVID-19: an Italian local register-based study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-051506.R1
Article Type:	Original research
Date Submitted by the Author:	11-Jul-2021
Complete List of Authors:	Fortunato, Francesca; University of Foggia, Department of Medical and Surgical Sciences; Foggia University Hospital, Department of Hygiene Martinelli, Domenico; University of Foggia, Department of Medical and Surgical Sciences; Foggia University Hospital, Department of Hygiene Lo Caputo, Sergio; University of Foggia, Department of Clinical and Experimental Medicine; Foggia University Hospital, Infectious Diseases Unit Santantonio, Teresa; University of Foggia, Department of Clinical and Experimental Medicine; Foggia University Hospital, Infectious Diseases Unit Dattoli, Vitangelo; Foggia University Hospital Lopalco, Pier Luigi; University of Salento, Department of Biological and Environmental Sciences and Technology Prato, Rosa; University of Foggia, Department of Medical and Surgical Sciences; Foggia University Hospital, Department of Hygiene
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Infectious diseases
Keywords:	COVID-19, Epidemiology < INFECTIOUS DISEASES, Public health < INFECTIOUS DISEASES, PUBLIC HEALTH, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Sex and gender differences in COVID-19: an Italian local register-based study

Francesca Fortunato¹, Domenico Martinelli¹, Sergio Lo Caputo², Teresa Santantonio², Vitangelo Dattoli³, Pier Luigi Lopalco⁴, and Rosa Prato^{1*}

¹ Department of Medical and Surgical Sciences, University of Foggia; Department of Hygiene, Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; francesca.fortunato@unifg.it, domenico.martinelli@unifg.it

² Department of Clinical and Experimental Medicine, University of Foggia; Infectious Diseases Unit, Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; sergio.locaputo@unifg.it, teresa.santantonio@unifg.it

³ Policlinico Riuniti University Hospital of Foggia, Foggia, Italy.; vitangelodattoli@gmail.com

⁴ Department of Biological and Environmental Sciences and Technology, University of Salento, Lecce, Italy; p.lopalco@regione.puglia.it

*Correspondence: rosa.prato@unifg.it; Ospedale "Colonnello D'Avanzo", Viale degli Aviatori, 2 - 71122 Foggia, Italy; Tel.: +39-0881-733091 (R.P.)

Word count: 2,853

Keywords: COVID-19; hospitalisation; virus clearance; death; sex differences; gender differences; Italy

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives

The present study aimed to explore differences in COVID-19 outcomes between male and female cases in the Apulian District of Foggia, Italy.

Design and setting

We performed a retrospective epidemiological study among all COVID-19 confirmed cases that occurred in the Apulian District of Foggia from February 29th to June 30th, 2020. The surveillance data from a regional registry (GIAVA-COVID©) were used.

Main outcomes

The main outcome measures were the proportion of hospitalisations, virus clearance and the case fatality rate.

Results

A total of 1,175 cases (50.7% female; median age: 55 years) were identified among 55,131 tests performed. The proportion of hospitalisation with COVID-19 diagnosis was 45.4% in men vs. 37.9% in women ($p<0.01$) while the average length of stay in hospitals was 31.3 ± 14.6 days in women vs. 26.8 ± 14.4 days in men ($p<0.01$). The proportion of cases who achieved virus clearance was higher in women (84.2%; days to clearance: 28.0 ± 12.1) than in men (79.3%; days to clearance: 29.4 ± 12.9 ; $p<0.05$). Men were associated with a significantly higher risk of dying from COVID-19 than women (case fatality rate [CFR] 16.1% vs. 10.4%; $p<0.01$). The mean time, from diagnosis to death was 14.5 ± 14.4 days in women compared with 10.6 ± 10.7 days in men ($p<0.01$). The male sex, age ≥ 55 years and presence of at least one underlying comorbidity significantly raised the risk of hospitalisation, persistent infection and death ($p<0.05$).

Conclusions

This study suggests that more attention should be paid to sex as a variable for the interpretation of COVID-19 data. Sex-disaggregated data will help clinicians to make appropriate patient-tailored medical decisions.

Strengths and Limitations of this Study

- This study provides sex-disaggregated data of COVID-19 cases at a district level, in Italy, contributing to a better understanding of who is being impacted the most by the pandemic and promoting a patient-tailored treatment approach.
- The robust methodology of the present study enabled to accurately correlate the case demographics with COVID-19 clinical response.
- The data related to the viral clearance, which reflect the diversified course of the disease according to the individual immune response, are confirmatory of sex difference in COVID-19.
- The data collected are highly homogeneous as they are strictly related to the first epidemic wave and provide an accurate picture of the impact of sex and age on COVID-19 outcomes in Italy during the initial phase of the pandemic.
- As the majority of the sex-disaggregated data available in the literature, the data presented in our study are not adjusted for lifestyle, profession, social or behavioural differences.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel coronavirus causing the current pandemic, which has resulted in millions of infections and hundreds of thousands of deaths worldwide. As of March 10th 2021, a total of 3,069,625 cases of confirmed SARS-COV-2 have been reported in Italy with a case fatality rate (CFR) of 3.2%.^[1]

The clinical manifestations of SARS-CoV-2 vary from asymptomatic infection to severe or critical disease.^[2]

Older age and comorbidities such as hypertension, cardiovascular disease, diabetes, chronic respiratory disease are associated with severe disease and death.^[3–5] Sex and gender have been identified as additional risk factors contributing to heterogeneous COVID-19 outcomes.^[2] Indeed, several studies have reported sex bias in COVID-19 case fatalities. It is observed that men have a higher risk of developing a severe form of the disease compared with women, highlighting the importance of sex-disaggregated data of COVID-19 cases.^[6] The initial reports from China followed by data from several European countries have shown similar numbers of confirmed cases between men and women.^[7,8] However, the severity of COVID-19, measured as hospitalisation, admission to intensive care units (ICUs) and fatality rate, is 2-fold higher in men than women.^[8,9] Studies in China, South Korea, United States, United Kingdom and Italy have reported higher case fatality rates and worst disease outcomes in male cases than in female cases.^[7,10–14] In some of these studies, the higher fatality rate in men was observed even after adjusting for confounding factors such as age and comorbidities.^[7,14] Additionally, in Italy, the higher fatality rate in men (age range: 40-80 years) is confirmed when the healthcare worker population is selectively studied.^[1]

The reasons for the differences in COVID-19 outcome and progression between men and women remain unclear. On one hand, biological factors, such as chromosomal and hormonal differences between men and women, may influence their susceptibility to infections, immune responses and progression of the disease.^[9,15–17] On the other hand, gender-related factors including psychological, social and behavioural differences between men and women may affect SARS-CoV-2 exposure, presence of comorbidities, treatment initiation and compliance, and COVID-19 mortality.^[18,19]

In this study, we used the surveillance data from a regional registry containing all confirmed cases of COVID-19 that occurred in the Foggia District (Apulia region, Italy), as of late June 2020, after the

end of the first epidemic wave. We aimed to explore the sex differences in hospitalisation, virus clearance, and deaths.

METHODS

Study population and design

We conducted a retrospective epidemiological study among COVID-19 cases that occurred in the Foggia District, Apulia region, Italy, from February 29th to June 30th, 2020. Foggia District is the third-largest Apulian District, with an estimated population of 616,310 residents (51% women) as of January 1st 2020.[20]

We used the surveillance data from a regional registry (GIAVA – COVID©) which was developed based on the WHO Go.Data outbreak investigation tool to manage the emergency.[21] GIAVA – COVID© includes functionalities for investigation and follow-up of cases and contacts, contact tracing, laboratory and clinical data collection. The collected information includes age, sex, residence location, date of disease onset, date of diagnosis, date of hospital admission, date of COVID-19 test results (positive or negative), date of death, presence of underlying diseases, case outcomes (hospitalisation, virus clearance and death), and disease severity (mild, moderate, severe, or critical).[22] The disease classification was duly updated according to clinical evolution of each case.

This study included all laboratory-confirmed cases defined as any person meeting the laboratory criterion (i.e. detection of SARS-CoV-2 nucleic acid or antigen in a clinical specimen).[23]

The proportion of hospitalisation was defined as the proportion of infected individuals undergoing hospitalisation among the total number of infected individuals. The proportion of individuals who achieved virus clearance was defined as the proportion of clinically recovered individuals with laboratory evidence of viral RNA clearance from the upper respiratory tract (two serial negative PCR tests at least 24 hours apart) among the total number of infected individuals. The case fatality rate was defined as the proportion of deaths among the total number of confirmed cases.

Statistical Analysis

Categorical variables were summarised as counts and percentages in each category. Data for continuous variables were expressed as medians (interquartile ranges [IQR]) and means (\pm standard

deviation [SD]). Normality of data was tested by the Kolmogorov-Smirnov test. Differences in continuous variables were assessed with Student's t-test or Mann-Whitney U test, depending on whether continuous variables were normally distributed or not, respectively. Significant differences in categorical variables were assessed using the Chi-square test or Fisher's exact test when appropriate and the odds ratio (OR) with 95% confidence interval (CI). Multivariate logistic regression analysis was performed to evaluate whether demographics (sex: male vs. female; age group: above vs. below the median age) and clinical characteristics (presence vs. absence of at least one underlying medical condition) were independently associated with hospitalisation, virus clearance, and deaths. The analysis was conducted with STATA/SE 15.0.

RESULTS

Between February 29th and June 30th, 2020, a total of 1,175 cases (50.7% female; median age: 55 years, IQR: 40-71 years) were diagnosed with COVID-19 in the Foggia District, Apulia region, Italy. The female positivity rate was 2.02% among 29,475 tests performed and the male positivity rate was 2.25% among 25,656 tests performed (chi-square $p > 0.05$).

Comparison of demographics and clinical characteristics of men versus women are shown in Table 1. A total of 373 cases (31.7%) had underlying medical conditions, including cardiovascular disease (63.3%), diabetes (19.6%), chronic pulmonary disease (13.9%), cancer (10.7%), neurological diseases (9.9%), chronic kidney disease (9.4%), and obesity (with body mass index [BMI] between 30-40 kg/m² or higher) (6.7%). Nearly 50% of cases were asymptomatic or with mild disease, 14.4% had moderate disease, 20.9% developed a severe disease and 3.2% progressed to a critical stage. There was no significant difference in age, underlying comorbidities (with the exception of diabetes), and disease severity distributions between the male and female groups (Table 1).

Table 1. Comparison of characteristics between male and female COVID-19 cases in the Foggia District, Apulia region, Italy

Characteristics	Male	Female	Total	OR (95% CI)	χ^2	p value
No. of cases (%)	579 (49.3)	596 (50.7)	1,175			
Median age (IQR), years	56 (40-70)	54.5 (38-74)	55 (40-71)			
Mean age (\pmSD), years	54.3 \pm 21.1	54.5 \pm 22.6	54.4 \pm 21.8			0.4291
Age group, No. (%):						
0-9	14 (2.4)	16 (2.7)	30 (2.6)	Ref.		
10-19	23 (4.0)	22 (3.7)	45 (3.8)	1.19 (0.43-3.34)	0.1	0.7061
20-29	43 (7.4)	53 (8.9)	96 (8.2)	0.93 (0.38-2.30)	0.03	0.8571
30-39	57 (9.8)	64 (10.7)	121 (10.3)	1.02 (0.42-2.47)	0.00	0.9655
40-49	91 (15.7)	95 (15.9)	186 (15.8)	1.09 (0.47-2.57)	0.05	0.8184
50-59	105 (18.1)	108 (18.1)	213 (18.1)	1.11 (0.48-2.59)	0.07	0.7874
60-69	96 (16.6)	64 (10.7)	160 (13.6)	1.71 (0.72-4.07)	1.84	0.1747
70-79	71 (12.3)	69 (11.6)	140 (11.9)	1.17 (0.49-2.81)	0.16	0.6874
80-89	64 (11.1)	72 (12.1)	136 (11.6)	1.01 (0.43-2.44)	0.00	0.9689
≥ 90	15 (2.6)	33 (5.5)	48 (4.1)	0.52 (0.18-1.48)	1.88	0.1705
Comorbidity, No (%)						
None	388 (67.0)	414 (69.5)	802 (68.3)	Ref.		
At least one comorbidity	191 (33.0)	182 (30.5)	373 (31.7)	1.1 (0.86-1.43)	0.75	0.3860
Cardiovascular disease	126 (66.9)	110 (60.4)	236 (63.3)	1.27 (0.81-1.97)	1.23	0.2682
Diabetes	49 (25.7)	24 (13.2)	73 (19.6)	2.27 (1.29-4.01)	9.20	0.0024
Chronic pulmonary disease	30 (15.7)	22 (12.1)	52 (13.9)	1.35 (0.72-2.58)	1.02	0.3132
Cancer	23 (12.0)	17 (9.3)	40 (10.7)	1.32 (0.65-2.75)	0.71	0.3994
Neurological diseases	15 (7.9)	22 (12.1)	37 (9.9)	0.62 (0.29-1.30)	1.87	0.1715
Chronic kidney disease	22 (11.5)	13 (7.1)	35 (9.4)	1.69 (0.78-3.78)	2.10	0.1475
Obesity	13 (6.8)	12 (6.6)	25 (6.7)	1.03 (0.42-2.55)	0.01	0.9345
Other metabolic diseases	5 (2.6)	10 (5.5)	15 (4.0)	0.46 (0.12-1.52)	2.00	0.1575
Liver disease	10 (5.2)	4 (2.2)	14 (3.8)	2.45 (0.69-10.91)	2.38	0.1228
Disease Severity, No. (%)						
Asymptomatic	92 (15.9)	113 (19.0)	205 (17.4)	Ref.		
Critical	23 (4.0)	15 (2.5)	38 (3.2)	1.88 (0.88-4.11)	3.15	0.0760
Severe	126 (21.8)	120 (20.1)	246 (20.9)	1.28 (0.87-1.90)	1.80	0.1796
Moderate	80 (13.8)	89 (14.9)	169 (14.4)	1.10 (0.72-1.69)	0.23	0.6348
Mild	165 (28.5)	197 (33.1)	362 (30.8)	1.02 (0.72-1.47)	0.03	0.8718

CI: confidence interval; IQR: interquartile range; OR: odds ratio; Ref.: reference group; SD: standard deviation

The proportion of hospitalisation among COVID-19 cases was estimated to be 41.6%, with a significant difference observed between men (45.4%) and women (37.9%; $p<0.01$). While the average length of stay in hospitals was significantly higher in women (31.3 ± 14.6 days) than in men (26.8 ± 14.4 days; $p<0.01$), there were more women aged ≥ 55 years hospitalised ($p<0.01$). The proportion of cases who achieved virus clearance was 82%, higher in women (84.2%; days to clearance: 28.0 ± 12.1) than in men (79.3%; days to clearance: 29.4 ± 12.9 ; $p<0.05$). A total of 155 deaths occurred among all cases for an overall CFR of 13.2%. Men were associated with a significantly higher risk of dying from COVID-19 than women (16.1% vs. 10.4%; $p<0.01$). The mean time, from diagnosis to death was higher in women (14.5 ± 14.4 days) compared with men (10.6 ± 10.7 days; $p<0.01$) (Table 2).

Table 2. Comparison of outcomes between male and female COVID-19 cases in the Foggia District, Apulia region, Italy

Characteristics	Male	Female	OR (95% CI)	χ^2	p value
Hospitalisation, No. (%)	263 (45.4)	226 (37.9)	1.36 (1.07-1.73)	6.81	0.0091
Mean age (\pm SD), years	66.2 \pm 16.0	70.2 \pm 18.8			0.0053
Mean length-of-stay in hospital (\pm SD), days	26.8 \pm 14.4	31.3 \pm 14.6			0.0032
Median length-of-stay in hospital (IQR), days	24 (17-35)	29 (19-41)			
Virus clearance (yes), No. (%)	459 (79.3)	502 (84.2)	0.72 (0.53-0.97)	4.84	0.0278
Mean time-to-virus-clearance (\pm SD), days	29.4 \pm 12.9	28.0 \pm 12.1			0.0432
Median time-to-virus-clearance (IQR), days	25 (18-35)	27 (19-37)			
Deaths, No. (%)	93 (16.1)	62 (10.4)	1.65 (1.15-2.36)	8.21	0.0042
Mean time-to-death (\pm SD), days	10.6 \pm 10.7	14.5 \pm 14.4			0.0282
Median time-to-death (IQR), days	8 (3-16)	10 (4-23)			

CI: confidence interval; IQR: interquartile range; OR: odds ratio; SD: standard deviation

The male sex, age ≥ 55 years and underlying comorbidities (presence of at least a condition among those listed in Table 1) significantly raised the risk of hospitalisation, persistent infection and death ($p<0.05$; Table 3).

Table 3. Multivariate logistic regression analysis of demographics and clinical characteristics of COVID-19 cases in the Foggia District, Apulia region, Italy

Characteristics	Hospitalisation		Virus clearance (no)		Deaths	
	OR (95% CI)	p value	OR (95% CI)	p value	OR (95% CI)	p value
Sex (male vs. female)	1.52 (1.15-2.20)	0.003	1.51 (1.08-2.09)	0.014	2.33 (1.52-3.58)	0.000
Age group (≥55 years vs. <55 years)	1.83 (1.68-1.99)	0.000	1.62 (1.47-1.78)	0.000	2.62 (2.22-3.07)	0.000
At least one comorbidity (yes vs. no)	1.99 (1.47-2.69)	0.000	1.63 (1.16-2.29)	0.004	1.94 (1.28-2.93)	0.002

CI: confidence interval; OR: odds ratio

DISCUSSION

Our registry-based surveillance study of 1,175 COVID-19 cases, well characterised from both demographic and clinical points of view, highlighted a male bias in COVID-19 outcomes. Based on the herein presented data, men are more likely to be hospitalised than women and the proportion of male cases achieving virus clearance is lower compared with female cases. Furthermore, men require longer periods to achieve virus clearance, have a higher fatality rate and faster progression to death.

A male bias (male-to-female ratio >1.1) in COVID-19 mortality is currently reported in 75 of the 94 countries that have provided sex-disaggregated data (as of March 10th 2021). At the global level, a higher number of men are hospitalised or admitted to ICU compared with women.[24] Additionally, several studies have demonstrated that men with COVID-19 are at higher risk of death and severe form of infection than women.[25,26] A recent meta-analysis of 3,111,714 reported global cases demonstrated that, whilst there is no difference in the proportion of male and female COVID-19 cases, men have higher odds of death (OR=1.39; 95% CI=1.31,1.47) compared with women.[27]. Similarly, our study presents a comparable proportion of women and men with confirmed COVID-19 (50.7% vs. 49.3%) and similar rates of positivity for infection (2.02% vs. 2.25%, $p > 0.05$). Therefore, the observed differences cannot be attributed to a prevalence of COVID-19 in the male sex.

The sex distribution of confirmed cases observed in the Foggia District, Apulia region is in line with the overall sex distribution of cases observed in Italy and other European countries.[1,8,28] Although in the early phase of the pandemic in Italy a higher prevalence of COVID-19 was observed

1
2
3 194 in men compared with women, this disproportion became less evident with the progression of the
4
5 195 pandemic. This variability may be explained by the different surveillance approaches adopted during
6
7 196 the pandemic since a symptom-based screening led to an underestimation of asymptomatic to mild
8
9 197 cases during the first epidemic wave. In Italy, after the end of the first epidemic wave (June 30th
10
11 198 2020), a higher number of male cases was observed in the 0-9, 10-19, 60-69 and 70-79 years age
12
13 199 groups (52.7%, 50.1%, 59.5%, 57.1%, respectively) compared with female cases, whereas a nearly
14
15 200 4-times higher number of female cases was observed in the >90 years age group.[29] On the
16
17 201 contrary, as of March 10th 2021, the number of confirmed COVID-19 cases is slightly higher in
18
19 202 women both in the overall Italian territory (51.4% in women vs. 48.6% in men) and in Apulia (51.%
20
21 203 in women vs. 48.4% in men).[1,30]
22
23 204 Differences in disease incidence, morbidity and mortality between sexes have also been observed
24
25 205 in other infectious diseases such as the severe acute respiratory syndrome coronavirus (SARS-COV-
26
27 206 1) and the Middle East respiratory syndrome coronavirus (MERS-COV) with men being more
28
29 207 susceptible than women to the infection and having a worse outcome.[31,32] The difference in
30
31 208 mortality between men and women suggests that women are either less prone to develop severe
32
33 209 complications or are less likely to die because of severe complications.[33]
34
35 210 The reasons behind these sex-related differences are probably pathogen-specific and of
36
37 211 multifactorial origin.[26] The three main determinants so far proposed to explain male-female
38
39 212 disparities in SARS-CoV-2 infection are differences in immune function associated with the X
40
41 213 chromosome, the effects of sex hormones, gender-related behavioural and socio-cultural
42
43 214 differences.[2,6,16,17] For example, the localisation of angiotensin-converting enzyme-2 (ACE2)
44
45 215 and Toll-like receptor 7 (TLR7) genes in the X chromosome and the mono-allelic versus the bi-allelic
46
47 216 presence may help explain the increased risk of COVID-19 for males compared with females.[34]
48
49 217 From a biological point of view, women seem to have a stronger immune system, weaker cytokine-
50
51 218 based pro-inflammatory response and lower levels of ACE2, an essential component for the
52
53 219 entrance of COVID-19 into the cells.[2,16,35–37] In this context, oestrogens seem to play a key
54
55 220 protective role. Oestrogen levels vary with age, rising in prepubertal individuals and decreasing with
56
57 221 age. Thus, the age-associated decline in oestradiol levels might be an explanation for the higher
58
59 222 susceptibility and severe progression of COVID-19 in older subjects.[38]
60
223 Our study highlights that, alongside sex, age and comorbidity are risk factors increasing
224 hospitalisation and death, and decreasing virus clearance. That COVID-19 severity increases with

age became evident since the beginning of the pandemic. Early studies from China and Italy showed that older age was associated with a greater risk of developing acute respiratory distress syndrome (ARDS), severe lung disease and death.[5,10] A recent meta-analysis of 55 studies and 10,014 COVID-19 cases confirmed that older age (≥ 50 years), together with comorbidities, significantly affects the prognosis and severity of COVID-19.[3] A further study investigated whether male bias in COVID-19 mortality was maintained at every age. It analysed data collated by the National Institute for Demographic Studies from national statistical agencies across England and Wales, France, Germany, Italy, Netherlands, Portugal, Korea, and Spain, including a population of 194,349,591 men and 201,715,364 women from the beginning of the pandemic until June 21st 2020. The overall male-to-female mortality sex ratio per 100,000 population was 1.4 (crude ratio 1.3). This ratio varied with age: 0.81 for subjects aged 0-9 years; 1.9 in the 40-49 year age group, 2.3 in the 50-59 year age group; 2.6 in the 60-69 year age group and 1.65 in subjects older than 80 years.[39] How the male vs. female difference in mortality, hospitalisation and virus clearance progresses with age is an aspect that warrants further investigation. In this context, stratification of the sex-disaggregated data provided in our study by age group could be relevant to better understand to what extent women are genetically protected from COVID-19. Interestingly, in our study, the stratification of the population by a cut-off age of 55 years highlighted a higher hospitalisation rate in the subgroup of women aged ≥ 55 years, suggesting the role of the reduction of hormonal protection with age.

One of the main hypotheses that have been postulated to justify the observed sex heterogeneity in the immune response to SARS-CoV-2 infection is the different genetic profile. Increasing evidence from patient populations highlights a substantial contribution of human genetic factors to the diversified susceptibility to SARS-CoV-2 infection and/or COVID-19 severity. In this context, a differential response to COVID-19 has also been observed among individuals with ethnicity-based differences in their genetic profile.[34] For instance, the distribution of the gene cluster on chromosome 3, that has been recently identified as the major genetic risk factor for severe COVID-19, differs among populations of different ethnic background (i.e. Asian, European and African populations).[40]

Lastly, gender-related differences in lifestyles and social roles require careful considerations as they are believed to greatly influence the onset, course and outcome of COVID-19. It has been proposed that smoking and alcohol consumption, alongside poor eating habits, more frequently found in men than women, may lead to a higher incidence of comorbidities in men compared with women

1
2
3 257 explaining the higher male mortality observed on a global level.[18,41] However, it must be noted
4
5 258 that no significant difference in underlying comorbidities (except for diabetes) between men and
6
7 259 women was found in our study. There may be other behavioural and social differences favouring
8
9 260 women as men are more reluctant to follow hand hygiene and seek preventive care.[42] On the
10
11 261 other hand, women might be more easily exposed to SARS-CoV-2 infection in both professional and
12
13 262 household settings. Indeed, women represent 70% of the health and social care workforce and more
14
15 263 often care for household members with COVID-19.[18,41]
16
17 264 The present study aimed to explore the differences in hospitalisation and death between men and
18
19 265 women at the local level taking into consideration COVID-19 confirmed cases in the Apulian District
20
21 266 of Foggia. The results are in line with what observed on a national and global level. Hospitalisation
22
23 267 and death are hard outcomes for monitoring the course and severity of the disease. Furthermore,
24
25 268 sex difference in virus clearance represents an added-value outcome of our study as it expresses
26
27 269 the immune response of the host.
28
29 270 However, it should not be neglected that one of the main limitations of our study is that the
30
31 271 presented data are not adjusted for lifestyle, profession, social or behavioural differences, all
32
33 272 relevant factors that could change the interpretation of the data and could further emphasise the
34
35 273 male-bias in COVID-19 severity and fatality. This limitation is a common feature of the majority of
36
37 274 sex-disaggregated data currently available. Indeed, due to practicability and ethical reasons, no
38
39 275 prospective study comparing an equal number of men and women under equal conditions of viral
40
41 276 exposure has been conducted to date. Therefore, we highlight the need of taking into account the
42
43 277 social, familiar and professional roles, alongside biological variables, in order to fully understand the
44
45 278 differences in COVID-19 outcome between men and women.
46
47 279 The main strength of our study consists in its robust methodology, which enabled an accurate
48
49 280 evaluation of the correlation between the case demographics (especially gender) and COVID-19
50
51 281 clinical response. Specifically, the collection of viral clearance data highlights a statistically
52
53 282 significant male-to-female difference and provides a plausible explanation for the observed
54
55 283 diversified course of the disease. Furthermore, the data collected in our study are highly
56
57 284 homogeneous as they are strictly related to the first epidemic wave, and provide an accurate picture
58
59 285 of the impact of sex and age on SARS-CoV-2 infection response in Italy during the initial phase of the
60
286 pandemic. The ethnic composition of the population included in our study is also highly
287 homogeneous and likely well representative of the Italian population or other Mediterranean
288 European populations.

CONCLUSIONS

Despite a comparable incidence of COVID-19 among the two sexes, a male bias in COVID-19 mortality is observed in the majority of the countries with available sex-disaggregated data. Our study provides sex-disaggregated data for the COVID-19 cases of the Apulian district of Foggia, Italy. It demonstrates that male sex, alongside older age (age ≥ 55 years) and presence of at least one comorbidity, is associated with a greater risk of hospitalisation and death, and lower virus clearance. Therefore, more attention should be paid to sex as a variable for the interpretation of COVID-19 data. This study will help clinicians to make appropriate patient-tailored medical decisions based on patient sex, age and comorbidities. Future investigations providing data adjusted for gender-related factors (social, familial and professional roles) are warranted.

Acknowledgements The Authors would like to thank all the frontline health workers at Hospitals and Local Health Unit for their dedication and valuable work into pandemic control. A special thanks to Lucia Massi, Maria Rosa Valetto and Pietro Dri (Zadig, Scientific Publisher, Milan, Italy) for editorial assistance, manuscript development and writing support.

Contributors FF and RP conceptualised and designed the work, analysed and interpreted data, and wrote the manuscript. DM and PLL supervised the study, coordinated regional data collection and provided statistical support. SLC, TS and VD interpreted the results and critically reviewed the advanced version of the manuscript. All authors approved the final draft of the manuscript. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for profit sectors.

Competing interests The authors declare no competing interests.

Patient and public involvement Patients and/or the public were not involved either in the study design, conduct, reporting or in the dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval

The study was conducted in accordance with the principles expressed in the Declaration of Helsinki 1975, as revised in 2008. As this study constituted public health surveillance, ethical approval from institutional review board was not required. All data were provided and analysed anonymously.

Data availability statement No additional data available.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

For peer review only

REFERENCES

- 1 COVID-19 integrated surveillance data in Italy.
<https://www.epicentro.iss.it/en/coronavirus/sars-cov-2-dashboard> (accessed 10 March 2021).
- 2 Falahi S, Kenarkoohi A. Sex and gender differences in the outcome of patients with COVID-19. *J Med Virol* 2021;**93**:151–2.
- 3 Barek MA, Aziz MA, Islam MS. Impact of age, sex, comorbidities and clinical symptoms on the severity of COVID-19 cases: A meta-analysis with 55 studies and 10014 cases. *Heliyon* 2020;**6**:e05684.
- 4 Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA* 2020;**18**:1775–6.
- 5 Wu C, Chen X, Cai Y, *et al.* Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med* 2020;**180**:934.
- 6 Dana PM, Sadoughi F, Hallajzadeh J, *et al.* An Insight into the Sex Differences in COVID-19 Patients: What are the Possible Causes? *Prehosp Disaster Med* 2020;**4**:438–41.
- 7 Chen J, Bai H, Liu J, *et al.* Distinct Clinical Characteristics and Risk Factors for Mortality in Female Inpatients With Coronavirus Disease 2019 (COVID-19): A Sex-stratified, Large-scale Cohort Study in Wuhan, China. *Clin Infect Dis* 2020;**71**:3188–95.
- 8 Gebhard C, Regitz-Zagrosek V, Neuhauser HK, *et al.* Impact of sex and gender on COVID-19 outcomes in Europe. *Biol Sex Differ* 2020;**11**:29.
- 9 Klein SL, Dhakal S, Ursin RL, *et al.* Biological sex impacts COVID-19 outcomes. *PLOS Pathog* 2020;**16**:e1008570.
- 10 Borghesi A, Zigliani A, Masciullo R, *et al.* Radiographic severity index in COVID-19 pneumonia: relationship to age and sex in 783 Italian patients. *Radiol Med* 2020;**125**:461–4.
- 11 Chen T, Wu D, Chen H, *et al.* Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;**368**:m1091.
- 12 Dudley JP, Lee NT. Disparities in Age-Specific Morbidity and Mortality from SARS-CoV-2 in China and the Republic of Korea. *Clin Infect Dis* 2020;**71**:863–5.
- 13 Gavin W, Campbell E, Zaidi S-A, *et al.* Clinical characteristics, outcomes and prognosticators in adult patients hospitalized with COVID-19. *Am J Infect Control* 2021;**49**:158–65.
- 14 Williamson EJ, Walker AJ, Bhaskaran K, *et al.* Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;**584**:430–6.
- 15 Dana PM, Sadoughi F, Hallajzadeh J, *et al.* An Insight into the Sex Differences in COVID-19 Patients: What are the Possible Causes? *Prehospital and Disaster Medicine*. 2020;**35**:438–41.

1
2
3 359 16 Bienvenu LA, Noonan J, Wang X, *et al.* Higher mortality of COVID-19 in males: sex differences
4 360 in immune response and cardiovascular comorbidities. *Cardiovasc Res* 2020;**116**:2197–206.
5
6
7 361 17 Scully EP, Haverfield J, Ursin RL, *et al.* Considering how biological sex impacts immune
8 362 responses and COVID-19 outcomes. *Nat Rev Immunol* 2020;**20**:442–7.
9
10 363 18 Cataldo Claudia, Masella Roberta. Gender-related sociocultural differences and COVID-19:
11 364 what influence on the effects of the pandemic? *Epidemiol Prev* 2020;**44**:Suppl 2:398-399.
12
13
14 365 19 Griffith DM, Sharma G, Holliday CS, *et al.* Peer Reviewed: Men and COVID-19: A
15 366 Biopsychosocial Approach to Understanding Sex Differences in Mortality and
16 367 Recommendations for Practice and Policy Interventions. *Prev Chronic Dis* 2020;**17**:E63.
17
18 368 20 Statistiche demografiche ISTAT. <http://demo.istat.it/pop2020/index.html> (accessed 1 Feb
19 369 2021).
20
21
22 370 21 Go.Data: Managing complex data in outbreaks. <https://www.who.int/godata> (accessed 1 Feb
23 371 2021).
24
25 372 22 Clinical management of COVID-19. [https://www.who.int/publications/i/item/clinical-](https://www.who.int/publications/i/item/clinical-management-of-covid-19)
26 373 [management-of-covid-19](https://www.who.int/publications/i/item/clinical-management-of-covid-19) (accessed 10 March 2021).
27
28
29 374 23 Case definition for coronavirus disease 2019 (COVID-19), as of 3 December 2020.
30 375 <https://www.ecdc.europa.eu/en/covid-19/surveillance/case-definition> (accessed 1 Feb 2021).
31
32 376 24 Global Health 50/50. [https://globalhealth5050.org/the-sex-gender-and-covid-19-](https://globalhealth5050.org/the-sex-gender-and-covid-19-project/dataset/)
33 377 [project/dataset/](https://globalhealth5050.org/the-sex-gender-and-covid-19-project/dataset/) (accessed 10 March 2021).
34
35 378 25 Kragholm K, Andersen MP, Gerds TA, *et al.* Association between male sex and outcomes of
36 379 Coronavirus Disease 2019 (Covid-19) – a Danish nationwide, register-based study. *Clin Infect*
37 380 *Dis*;ciaa**924**.
38
39
40 381 26 Lakbar I, Luque-Paz D, Mege J-L, *et al.* COVID-19 gender susceptibility and outcomes: A
41 382 systematic review. *PLOS ONE* 2020;**15**:e0241827.
42
43
44 383 27 Peckham H, Gruijter NM de, Raine C, *et al.* Male sex identified by global COVID-19 meta-
45 384 analysis as a risk factor for death and ITU admission. *Nat Commun* 2020;**11**:1–10.
46
47 385 28 European Centre for Disease Prevention and Control (ECDC). COVID-19 situation board.
48 386 [https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-](https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-transmission-tab)
49 387 [transmission-tab](https://qap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html#subnational-transmission-tab) (accessed 10 March 2021).
50
51
52 388 29 Bollettino-sorveglianza-integrata-COVID-19_30 giugno 2020.
53 389 [https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-come-integrata-](https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-come-integrata-COVID-19_30-giugno-2020.pdf)
54 390 [COVID-19_30-giugno-2020.pdf](https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-come-integrata-COVID-19_30-giugno-2020.pdf) (accessed 10 March 2021).
55
56 391 30 Bollettino Epidemiologico Regione Puglia. Epidemia COVID-19, 10 marzo 2021.
57 392 https://www.regione.puglia.it/documents/65725/216593/Bollettino+Covid_10032021.pdf/6cf
58 393 [c1d87-a820-25c9-9832-2585c0f50099?t=1615382755715](https://www.regione.puglia.it/documents/65725/216593/Bollettino+Covid_10032021.pdf/6cf) (accessed 10 March 2021).
59
60

- 31 Karlberg J, Chong DSY, Lai WYY. Do men have a higher case fatality rate of severe acute respiratory syndrome than women do? *Am J Epidemiol* 2004;**159**:229–31.
- 32 Alghamdi IG, Hussain II, Almalki SS, *et al.* The pattern of Middle East respiratory syndrome coronavirus in Saudi Arabia: a descriptive epidemiological analysis of data from the Saudi Ministry of Health. *Int J Gen Med* 2014;**7**:417–23.
- 33 Penna C, Mercurio V, Tocchetti CG, *et al.* Sex-related differences in COVID-19 lethality. *Br J Pharmacol* 2020;**177**:4375–85.
- 34 Anastassopoulou C, Gkizarioti Z, Patrinos GP, *et al.* Human genetic factors associated with susceptibility to SARS-CoV-2 infection and COVID-19 disease severity. *Human Genomics* 2020;**14**:40.
- 35 Haitao T, Vermunt JV, Abeykoon J, *et al.* COVID-19 and Sex Differences: Mechanisms and Biomarkers. *Mayo Clin Proc* 2020;**95**:2189–203.
- 36 Takahashi T, Ellingson MK, Wong P, *et al.* Sex differences in immune responses that underlie COVID-19 disease outcomes. *Nature* 2020;**588**:315–20.
- 37 Takahashi T, Iwasaki A. Sex differences in immune responses. *Science* 2021;**371**:347–8.
- 38 Liang X. Is COVID-19 more severe in older men? *Postgrad Med J* 2020;**96**:426.
- 39 Bhopal SS, Bhopal R. Sex differential in COVID-19 mortality varies markedly by age. *The Lancet* 2020;**396**:532–3.
- 40 Zeberg H, Pääbo S. The major genetic risk factor for severe COVID-19 is inherited from Neanderthals. *Nature* 2020;**587**:610–2.
- 41 Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. *The Lancet* 2020;**395**:846–8.
- 42 Sharma G, Volgman AS, Michos ED. Sex Differences in Mortality From COVID-19 Pandemic: Are Men Vulnerable and Women Protected? *JACC Case Rep* 2020;**2**:1407–10.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—checklist of items that should be included in reports of observational studies

Sex and gender differences in COVID-19: an Italian local register-based study
Francesca Fortunato, Domenico Martinelli, Sergio Lo Caputo, Teresa Santantonio, Vitangelo Dattoli, Pier Luigi Lopalco, and Rosa Prato.

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1 (title) 2 (abstract)
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2 (abstract)
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4 (lines 65-92)
Objectives	3	State specific objectives, including any prespecified hypotheses	4-5 (lines 92,96)
Methods			
Study design	4	Present key elements of study design early in the paper	5 (lines 100-103)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5 (lines 100-112)
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	(a) Cross-sectional 5 (lines 115-120)
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5 (lines 103-107)
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5 (lines 104-107; 115-132)
Bias	9	Describe any efforts to address potential sources of bias	5-6 (lines 128-132)
Study size	10	Explain how the study size was arrived at	5-6 (lines 113-114)
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	5-6 (lines 121-132)

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	5-6 (lines 121-132)
		(b) Describe any methods used to examine subgroups and interactions	5-6 (lines 121-132)
		(c) Explain how missing data were addressed	Not applicable
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	
		(e) Describe any sensitivity analyses	Not applicable
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6 (lines 138-149) Table 1
		(b) Give reasons for non-participation at each stage	Not applicable
		(c) Consider use of a flow diagram	Not applicable
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6-7 (lines 132-140) Table 1
		(b) Indicate number of participants with missing data for each variable of interest	Not applicable
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	8-9 (lines 154-175) Tables 2, 3
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8-9 (lines 154-175) Tables 2, 3
		(b) Report category boundaries when continuous variables were categorized	Tables 2,3
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Tables 2,3
Discussion			
Key results	18	Summarise key results with reference to study objectives	9 (lines 177-182)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12 (lines 273-281)
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12 (lines 282-291)
Generalisability	21	Discuss the generalisability (external validity) of the study results	13 (lines 292-301)
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	13 (lines 312,313)

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.